

IJSPT
Clinical Viewpoint

Why Female Athletes Injure Their ACL's More Frequently? What can we do to mitigate their risk?

Holly Silvers-Granelli, PT, MPT, PhD
Velocity Physical Therapy, Santa Monica, CA
Major League Soccer Medical Research Committee, New York, New York

Contact information:

ORCID # 0000-0002-8959-9448
2716 Ocean Park Blvd., Suite 1065
Santa Monica, CA 90405
Tel: 310 591 8016
Email: Hollysilverspt@gmail.com

Anterior Cruciate Ligament (ACL) injuries are one of the most deleterious knee injuries reported in sport. They continue to confound the sports medicine community, particularly with respect to the high rates reported in girls and women.¹⁻⁴ There are approximately 200,000 to 250,000 ACL injuries that occur in the United States annually, a rate that has doubled over the last 20 years.^{5,6} Approximately 25% of these injuries occur in youth athletics; and this rate has been increasing by a rate of 2.5% annually in the United States⁷ and has increased by 147.8% over a 10 year period in Victoria, Australia.⁸ Although the overall rate of ACL injury is higher in males, primarily due to greater opportunity(s) to participate in contact sports, the relative risk of ACL injury in women is 3 to 8 times greater than males.^{9,10} In the National Collegiate Athletic Association (NCAA), the rate of ACL injuries incurred by collegiate females is three times higher compared to men.¹ In high school aged athletes (13-18 years), there is approximately 1.6-fold greater rate of ACL tears in females,¹¹ and a multisport female athlete is estimated to have a nearly 10% risk of incurring an ACL injury during her high school or secondary school career.¹²

A concerted effort has been made over the last three decades to address the complexities of the “sequence of prevention”; to mitigate ACL injury risk by virtue of the implementation of validated injury prevention program (IPP) interventions.^{13,14} A vast majority of these IPP’s were designed specifically to address ACL injury in females.¹⁵⁻¹⁹ The targeted risk factors included anatomic, environmental, hormonal, genetic and biomechanical.^{20,21} These neuromuscular IPP training programs, that addressed the biomechanical deficiencies, included in-season elements of strength, plyometrics, sport specific agility drills, proper landing technique, proprioception, proximal control and a biomechanical emphasis on addressing the most common

pathokinematic movement patterns associated with ACL injury mechanisms.^{22 23 24} Additional components of successful ACL IPPs include socio-economic feasibility, sports-specificity, physiological competency, neurocognitive and psychological (improving confidence and reducing fear) and implementation strategies that may improve overall program adoption.²⁵⁻²⁸ The IPP's were typically designed as in-season dynamic warm-up programs, included both intrinsic and extrinsic cues, were strategically offered at no or low-cost, included coaching and athlete web-based educational tools, and were time efficient to promote overall team compliance, program fidelity and adherence.^{19,29-31} The IPP's have been largely successful; reporting overall ACL injury reduction rates between 55-88%.^{16,17,19,31-33} The programs are designed to be introduced during preseason and continue throughout the season to mitigate biomechanical recidivism.^{34,35} Additionally, the day in which the IPP was performed resulted in even lower ACL injury rates, suggesting that a transient, neural preparedness and cortical control element may be favorably impacting overall biomechanics and motor control.^{30 36} Including principals of motor learning theories as a component of rehabilitation and in IPPs has led to improvements in efficiency of the motor cortex, ostensibly allowing the athlete to make improvements to their biomechanics while allowing them to interpret and process rapidly changing environmental stimuli due to improved neurocognitive availability.^{37,38} Optimization of IPPs must include a synergy of cognitive, perceptual, and motor processes to enhance the athletes' ability to respond to sport-specific demands with comprehensive and low-risk biomechanical movement strategies.³⁹

Recent studies have retrospectively analyzed injury mechanisms in male and female athletes to further elucidate the biomechanical pathokinematics specifically involved in the mechanism of injury.^{40 41,42 43 44-50} Video analysis of ACL injuries in male and female athletes have begun to effectively delineate high risk positioning associated with the injury, namely defensive and unanticipated play, with the injured player demonstrating at or near full hip and knee extension, perturbation to the trunk resulting in lateral trunk displacement, hip adduction and internal rotation, knee valgus, and tibial torsion.^{51 40 52} Females were more likely to be defending or in an unanticipated/reactive position and were more likely to tear their non-dominant limb.^{41,53}

Studies analyzing the role of peripheral fatigue and its' role in ACL injury have been in consistent⁵⁴. A study analyzing female ACL injury mechanisms suggested that fatigue was not correlated with injury, as 64% of injuries occurred in the first 30 minutes of a soccer match.^{49 55} However, peripheral fatigue has been shown to be a variable for women in Irish Amateur Rugby⁵⁶, altering biomechanics during landing performance at initial contact^{57 58}, increasing trunk flexion⁵⁹, and reducing peak knee extensor torque^{60 61}. Inclusion of fatigue as one of the metrics for IPP efficacy should be considered.⁵⁸ There is inherent complexity to determining the external validity of fatigue on ACL injury incidence. As fatigue increases, psychological stress may increase (stress, emotional lability) and physical response may decrease (performance, velocity, neuromuscular workload and intensity). The decrease in player intensity, performance, and velocity may be more reflective of central fatigue and may ultimately mitigate the overall risk of ACL injury.⁵⁸ The continued identification and

understanding of the intrinsic and extrinsic sex related ACL injury risk factors will increase the clinician's ability to elucidate and improve IPPs to effectively decrease the ACL injury rate in sport.

One of the major difficulties researchers are enduring, from a public health perspective, is achieving widespread program adoption and implementation of the established and validated IPP's. Despite the earnest efforts of researchers to mitigate ACL injury rate through the development and the evolution of the aforementioned IPPs, the programs' potential to reduce risk has been hindered by the overall low adoption rate of these programs. Interestingly, it has been well documented that high compliance to a scientifically vetted IPP can substantially mitigate ACL injury rates.⁶²⁻⁶⁷ Conversely, when overall compliance was low and the IPPs were performed less than once per week and/or with low program fidelity, the IPPs were found to be largely ineffective.⁶⁸⁻⁶⁹ Upon analyzing coaching decisions to consistently using an IPP program, researchers determined that it requires a detailed understanding of the unique implementation context, including exercise variety and modification to expand its' reach, sport specific exercises, incorporating sport specific equipment, time and cost efficacy, greater exercise variations and increased difficulty in program progressions.^{70,71} These alterations should be heavily considered in IPP design, as the cohesive and consistent implementation of IPPs is a very viable, impactful, and cost-effective option to reducing the overall rate of ACL injury.⁷² Several studies have demonstrated a positive effect of IPPs on its effectiveness as a warm-up and overall athlete performance.⁷³⁻⁷⁶ Optimizing implementation and team compliance, particularly at the youth and recreational levels, lies within the coaching decision making

paradigm. The notion of improved performance, recognized by and improved win-loss record, and decreasing overall injury rate to improve player availability may optimally incentivize coaches and players in incorporate an IPP with regularity. ⁷⁷ (Silvers-Granelli, in peer review, Sports Health).

A more nuanced narrative has recently emerged with respect to challenging the prevailing ACL injury prevention debate; are females truly more vulnerable to ACL injury or is this simply a consequence of a series of gendered societal and environmental decisions? Most ACL epidemiological and mechanism studies have been centered around gender-based biology, without considering other social, economic, contextual, and environmental factors. There is a significant disparity in training, coaching and competitive resources in female sports. Despite the advent of the Title IX Educational Amendment in 1972, which prohibited sex discrimination in any education program or activity receiving federal financial assistance in the United States, there is a incongruity in what females are afforded in competitive sporting environments. ⁷⁸ This includes, but is not limited to, decreased overall salaries for coaching and professional play, diminished access to exercise equipment and high quality and consistent rehabilitation, lower standards for coaching, medical staffing and strength and conditioning professional experience, and decreased access to childcare and maternity benefits during their professional careers. ⁷⁹ The impact of ACL injury and reconstruction have also differed in males versus females. Upon a two-year longitudinal analysis, females have demonstrated reticence in return to play activity, exhibited through behavioral self-modulation, by virtue of a decrease in vigorous activity, decreased triple hop distance, and a shift away from team sport participation to mitigate

secondary injury risk.⁸⁰ This concerted decision, to decrease overall secondary risk through behavioral modification, may be partly due to the fact that the risk:reward balance that exists for men is simply not a realistic option for most women. It would behoove the research community to consider additional possibilities to the existing “biological element” influences that currently dominate the prevailing ACL injury prevention algorithm.

As we embark upon our fourth decade on the ACL injury mitigation journey, perhaps we “pivot” and discuss how we effectively disseminate information in a way that encompasses the current social, economic and environmental sex differences across sport. If we recognize the current inequity, and scientifically modify our algorithms, our prevention outreach and interventions may be perceived more favorably and just might increase their overall efficacy. Let us all be prescient as we attempt to minimize the current gender gaps present across sport and respond accordingly.

REFERENCES

1. Agel J, Arendt EA, Bershadsky B. Anterior cruciate ligament injury in national collegiate athletic association basketball and soccer: a 13-year review. *The American journal of sports medicine*. 2005;33(4):524-530.
2. Agel J, Rockwood T, Klossner D. Collegiate ACL Injury Rates Across 15 Sports: National Collegiate Athletic Association Injury Surveillance System Data Update (2004-2005 Through 2012-2013). *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*. 2016;26(6):518-523.
3. Junge A, Dvorak J. Injuries in female football players in top-level international tournaments. *Br J Sports Med*. 2007;41 Suppl 1:i3-7.
4. Junge A, Dvorak J. Injury surveillance in the World Football Tournaments 1998-2012. *Br J Sports Med*. 2013;47(12):782-788.
5. Yu B, Garrett WE. Mechanisms of non-contact ACL injuries. *British journal of sports medicine*. 2007;41 Suppl 1(Suppl 1):i47-i51.
6. Albright JC CJ, Graf BK. *Knee and leg: soft tissue trauma*. Vol 6. Rosemont, IL.: American Academy of Orthopaedic Surgeons; 1999.
7. Beck NA, Lawrence JTR, Nordin JD, DeFor TA, Tompkins M. ACL Tears in School-Aged Children and Adolescents Over 20 Years. *Pediatrics*. 2017:e20161877.
8. Shaw L, Finch CF. Trends in Pediatric and Adolescent Anterior Cruciate Ligament Injuries in Victoria, Australia 2005-2015. *International journal of environmental research and public health*. 2017;14(6).
9. Montalvo AM, Schneider DK, Yut L, et al. "What's my risk of sustaining an ACL injury while playing sports?" A systematic review with meta-analysis. *British Journal of Sports Medicine*. 2019;53(16):1003-1012.
10. Ardern CL, Ekås G, Grindem H, et al. 2018 International Olympic Committee consensus statement on prevention, diagnosis and management of paediatric anterior cruciate ligament (ACL) injuries. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2018.
11. Gornitzky AL, Lott A, Yellin JL, Fabricant PD, Lawrence JT, Ganley TJ. Sport-Specific Yearly Risk and Incidence of Anterior Cruciate Ligament Tears in High School Athletes: A Systematic Review and Meta-analysis. *The American journal of sports medicine*. 2016;44(10):2716-2723.
12. Bram JT, Magee LC, Mehta NN, Patel NM, Ganley TJ. Anterior Cruciate Ligament Injury Incidence in Adolescent Athletes: A Systematic Review and Meta-analysis. *The American journal of sports medicine*. 2020:363546520959619.
13. van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports medicine (Auckland, NZ)*. 1992;14(2):82-99.
14. Bolling C, van Mechelen W, Pasman HR, Verhagen E. Context Matters: Revisiting the First Step of the 'Sequence of Prevention' of Sports Injuries. *Sports Medicine*. 2018;48(10):2227-2234.
15. Mandelbaum B, Silvers H, Watanabe D, et al. Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes - 2-year follow-up. *American Journal of Sports Medicine*. 2005;33(7):1003-1010.

16. Gilchrist J, Mandelbaum BR, Melancon H, et al. A randomized controlled trial to prevent noncontact anterior cruciate ligament injury in female collegiate soccer players. *The American journal of sports medicine*. 2008;36(8):1476-1483.
17. Myklebust G, Engebretsen L, Braekken IH, Skjolberg A, Olsen OE, Bahr R. Prevention of noncontact anterior cruciate ligament injuries in elite and adolescent female team handball athletes. *Instructional course lectures*. 2007;56:407-418.
18. Silvers HJ, Giza ER, Mandelbaum BR. Anterior cruciate ligament tear prevention in the female athlete. *Current sports medicine reports*. 2005;4(6):341-343.
19. Walden M, Atroshi I, Magnusson H, Wagner P, Hagglund M. Prevention of acute knee injuries in adolescent female football players: cluster randomised controlled trial. *BMJ (Clinical research ed)*. 2012;344:e3042.
20. Griffin LY, Agel J, Albohm MJ, et al. Noncontact anterior cruciate ligament injuries: risk factors and prevention strategies. *J Am Acad Orthop Surg*. 2000;8(3):141-150.
21. Griffin LY, Albohm MJ, Arendt EA, et al. Understanding and preventing noncontact anterior cruciate ligament injuries: a review of the Hunt Valley II meeting, January 2005. *The American journal of sports medicine*. 2006;34(9):1512-1532.
22. Bizzini M JA, Dvorak J. . Implementation of the FIFA 11+ football warm up program: how to approach and convince the Football associations to invest in prevention. *Br J Sports Med*. 2013;Aug;47(12):803-806.
23. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: Part 1, mechanisms and risk factors. *The American journal of sports medicine*. 2006;34(2):299-311.
24. Thorborg K, Krommes KK, Esteve E, Clausen MB, Bartels EM, Rathleff MS. Effect of specific exercise-based football injury prevention programmes on the overall injury rate in football: a systematic review and meta-analysis of the FIFA 11 and 11+ programmes. *Br J Sports Med*. 2017;51(7):562-571.
25. Bien DP. Rationale and implementation of anterior cruciate ligament injury prevention warm-up programs in female athletes. *J Strength Cond Res*. 2011;25(1):271-285.
26. Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. . A conceptual framework for implementation fidelity. *Implementation Science*. 2007;2(40).
27. Donaldson A, Finch CF. Applying implementation science to sports injury prevention. *Br J Sports Med*. 2013;47(8):473-475.
28. O'Brien J, Finch CF. The implementation of musculoskeletal injury-prevention exercise programmes in team ball sports: a systematic review employing the RE-AIM framework. *Sports medicine (Auckland, NZ)*. 2014;44(9):1305-1318.
29. Bizzini M, Dvorak J. FIFA 11+: an effective programme to prevent football injuries in various player groups worldwide-a narrative review. *Br J Sports Med*. 2015;49(9):577-579.
30. Silvers-Granelli H, Mandelbaum B, Adeniji O, et al. Efficacy of the FIFA 11+ Injury Prevention Program in the Collegiate Male Soccer Player. *The American journal of sports medicine*. 2015.
31. Soligard T, Myklebust G, Steffen K, et al. Comprehensive warm-up programme to prevent injuries in young female footballers: cluster randomised controlled trial. *BMJ (Clinical research ed)*. 2008;337:a2469.

32. Hagglund M, Walden M, Atroshi I. Preventing knee injuries in adolescent female football players - design of a cluster randomized controlled trial [NCT00894595]. *BMC musculoskeletal disorders*. 2009;10:75.
33. Hewett TE, Myer GD, Ford KR. Reducing knee and anterior cruciate ligament injuries among female athletes: a systematic review of neuromuscular training interventions. *The journal of knee surgery*. 2005;18(1):82-88.
34. Mandelbaum BR, Silvers HJ, Watanabe DS, et al. Effectiveness of a neuromuscular and proprioceptive training program in preventing anterior cruciate ligament injuries in female athletes: 2-year follow-up. *The American journal of sports medicine*. 2005;33(7):1003-1010.
35. Mouton C, Gokeler A, Urhausen A, Nührenbörger C, Seil R. High Incidence of Anterior Cruciate Ligament Injuries Within the First 2 Months of the Season in Amateur Team Ball Sports. *Sports health*. 2021:19417381211014140.
36. Grooms DR, Kiefer AW, Riley MA, et al. Brain-Behavior Mechanisms for the Transfer of Neuromuscular Training Adaptions to Simulated Sport: Initial Findings From the Train the Brain Project. *J Sport Rehabil*. 2018;27(5):1-5.
37. Gokeler A, Neuhaus D, Benjaminse A, Grooms DR, Baumeister J. Principles of Motor Learning to Support Neuroplasticity After ACL Injury: Implications for Optimizing Performance and Reducing Risk of Second ACL Injury. *Sports medicine (Auckland, NZ)*. 2019;49(6):853-865.
38. Grooms DR, Page SJ, Nichols-Larsen DS, Chaudhari AM, White SE, Onate JA. Neuroplasticity Associated With Anterior Cruciate Ligament Reconstruction. *The Journal of orthopaedic and sports physical therapy*. 2017;47(3):180-189.
39. Kiefer AW, Myer GD. Training the Antifragile Athlete: A Preliminary Analysis of Neuromuscular Training Effects on Muscle Activation Dynamics. *Nonlinear Dynamics Psychol Life Sci*. 2015;19(4):489-510.
40. Walden M, Krosshaug T, Bjerneboe J, Andersen TE, Faul O, Hagglund M. Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *British journal of sports medicine*. 2015;49(22):1452-1460.
41. Brophy RH, Stepan JG, Silvers HJ, Mandelbaum BR. Defending Puts the Anterior Cruciate Ligament at Risk During Soccer: A Gender-Based Analysis. *Sports health*. 2015;7(3):244-249.
42. Johnston JT, Mandelbaum BR, Schub D, et al. Video Analysis of Anterior Cruciate Ligament Tears in Professional American Football Athletes. *The American journal of sports medicine*. 2018;46(4):862-868.
43. Della Villa F, Buckthorpe M, Grassi A, et al. Systematic video analysis of ACL injuries in professional male football (soccer): injury mechanisms, situational patterns and biomechanics study on 134 consecutive cases. *Br J Sports Med*. 2020.
44. Della Villa F, Buckthorpe M, Grassi A, et al. Systematic video analysis of ACL injuries in professional male football (soccer): injury mechanisms, situational patterns and biomechanics study on 134 consecutive cases. *Br J Sports Med*. 2020;54(23):1423-1432.
45. Della Villa F, Di Paolo S, Santagati D, et al. A 2D video-analysis scoring system of 90° change of direction technique identifies football players with high knee abduction

- moment. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA*. 2021.
46. Dix C, Arundale A, Silvers-Granelli H, Marmon A, Zarzycki R, Snyder-Mackler L. BIOMECHANICAL MEASURES DURING TWO SPORT-SPECIFIC TASKS DIFFERENTIATE BETWEEN SOCCER PLAYERS WHO GO ON TO ANTERIOR CRUCIATE LIGAMENT INJURY AND THOSE WHO DO NOT: A PROSPECTIVE COHORT ANALYSIS. *International journal of sports physical therapy*. 2020;15(6):928-935.
 47. Grassi A, Tosarelli F, Agostinone P, Macchiarola L, Zaffagnini S, Della Villa F. Rapid Posterior Tibial Reduction After Noncontact Anterior Cruciate Ligament Rupture: Mechanism Description From a Video Analysis. *Sports health*. 2020;12(5):462-469.
 48. Larwa J, Stoy C, Chafetz RS, Boniello M, Franklin C. Stiff Landings, Core Stability, and Dynamic Knee Valgus: A Systematic Review on Documented Anterior Cruciate Ligament Ruptures in Male and Female Athletes. *International journal of environmental research and public health*. 2021;18(7).
 49. Lucarno S, Zago M, Buckthorpe M, et al. Systematic Video Analysis of Anterior Cruciate Ligament Injuries in Professional Female Soccer Players. *The American journal of sports medicine*. 2021:3635465211008169.
 50. Scarborough DM, Linderman SE, Cohen VA, Berkson EM, Eckert MM, Oh LS. Neuromuscular Control of Vertical Jumps in Female Adolescents. *Sports health*. 2019;11(4):343-349.
 51. Shimokochi Y, Shultz SJ. Mechanisms of noncontact anterior cruciate ligament injury. *J Athl Train*. 2008;43(4):396-408.
 52. Stuelcken MC, Mellifont DB, Gorman AD, Sayers MG. Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *Journal of sports sciences*. 2016;34(16):1516-1522.
 53. Brophy R, Silvers HJ, Gonzales T, Mandelbaum BR. Gender influences: the role of leg dominance in ACL injury among soccer players. *Br J Sports Med*. 2010;44(10):694-697.
 54. Barber-Westin SD, Noyes FR. Effect of Fatigue Protocols on Lower Limb Neuromuscular Function and Implications for Anterior Cruciate Ligament Injury Prevention Training: A Systematic Review. *The American journal of sports medicine*. 2017;45(14):3388-3396.
 55. Alsubaie SF, Abdelbasset WK, Alkathiry AA, et al. Anterior cruciate ligament injury patterns and their relationship to fatigue and physical fitness levels - a cross-sectional study. *Medicine (Baltimore)*. 2021;100(1):e24171.
 56. Yeomans C, Kenny IC, Cahalan R, et al. Injury Trends in Irish Amateur Rugby: An Epidemiological Comparison of Men and Women. *Sports health*. 2021:1941738121997145.
 57. Gokeler A, Eppinga P, Dijkstra PU, et al. Effect of fatigue on landing performance assessed with the landing error scoring system (less) in patients after ACL reconstruction. A pilot study. *International journal of sports physical therapy*. 2014;9(3):302-311.
 58. Benjaminse A, Webster KE, Kimp A, Meijer M, Gokeler A. Revised Approach to the Role of Fatigue in Anterior Cruciate Ligament Injury Prevention: A Systematic Review with Meta-Analyses. *Sports medicine (Auckland, NZ)*. 2019;49(4):565-586.

59. Lessi GC, Serrão FV. Effects of fatigue on lower limb, pelvis and trunk kinematics and lower limb muscle activity during single-leg landing after anterior cruciate ligament reconstruction. *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA*. 2017;25(8):2550-2558.
60. Greig M. The influence of soccer-specific fatigue on peak isokinetic torque production of the knee flexors and extensors. *The American journal of sports medicine*. 2008;36:1403-1409.
61. Jones RI, Ryan B, Todd AI. Muscle fatigue induced by a soccer match-play simulation in amateur Black South African players. *Journal of sports sciences*. 2015;33(12):1305-1311.
62. Hagglund M, Atroshi I, Wagner P, Walden M. Superior compliance with a neuromuscular training programme is associated with fewer ACL injuries and fewer acute knee injuries in female adolescent football players: secondary analysis of an RCT. *Br J Sports Med*. 2013;47(15):974-979.
63. Silvers-Granelli HJ, Bizzini M, Arundale A, Mandelbaum BR, Snyder-Mackler L. Higher compliance to a neuromuscular injury prevention program improves overall injury rate in male football players. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2018:1-9.
64. Soligard T, Nilstad A, Steffen K, et al. Compliance with a comprehensive warm-up programme to prevent injuries in youth football. *Br J Sports Med*. 2010;44(11):787-793.
65. Steffen K, Emery CA, Romiti M, et al. High adherence to a neuromuscular injury prevention programme (FIFA 11+) improves functional balance and reduces injury risk in Canadian youth female football players: a cluster randomised trial. *Br J Sports Med*. 2013;47(12):794-802.
66. Sugimoto D, Myer GD, Bush HM, Klugman MF, Medina McKeon JM, Hewett TE. Compliance with neuromuscular training and anterior cruciate ligament injury risk reduction in female athletes: a meta-analysis. *J Athl Train*. 2012;47(6):714-723.
67. van Reijen M, Vriend I, van Mechelen W, Finch CF, Verhagen EA. Compliance with Sport Injury Prevention Interventions in Randomised Controlled Trials: A Systematic Review. *Sports medicine (Auckland, NZ)*. 2016;46(8):1125-1139.
68. Steffen K, Myklebust G, Olsen OE, Holme I, Bahr R. Preventing injuries in female youth football—a cluster-randomized controlled trial. *Scandinavian journal of medicine & science in sports*. 2008;18(5):605-614.
69. Beynon BD, Vacek P, Tourville TW, et al. Implementation of the FIFA 11+ Injury Prevention Program by High School Athletic Teams Did Not Reduce Lower Extremity Injuries: Response. *The American journal of sports medicine*. 2020;48(6):Np36-np37.
70. O'Brien J, Finch CF. Injury prevention exercise programmes in professional youth soccer: understanding the perceptions of programme deliverers. *BMJ Open Sport Exer Med*. 2016;2(1):e000075-e000075.
71. O'Brien J, Young W, Finch CF. The use and modification of injury prevention exercises by professional youth soccer teams. *Scan J Med Sci Sport*. 2017;27(11):1337-1346.
72. Junge A, Lamprecht M, Stamm H, et al. Countrywide campaign to prevent soccer injuries in Swiss amateur players. *The American journal of sports medicine*. 2011;39(1):57-63.
73. Faude O, Rößler R, Junge A. Football injuries in children and adolescent players: are there clues for prevention? *Sports Med*. 2013;43(9):819-837.

74. Rössler R, Donath L, Bizzini M, Faude O. A new injury prevention programme for children's football—FIFA 11+ Kids—can improve motor performance: a cluster-randomised controlled trial. *J Sports Sci.* 2016;34(6):549-556.
75. Zarei M, Abbasi H, Daneshjoo A, et al. Long-term effects of the 11+ warm-up injury prevention programme on physical performance in adolescent male football players: a cluster-randomised controlled trial. *J Sports Sci.* 2018;36(21):2447-2454.
76. Impellizzeri FM, Bizzini M, Dvorak J, Pellegrini B, Schena F, Junge A. Physiological and performance responses to the FIFA 11+ (part 2): a randomised controlled trial on the training effects. *Journal of sports sciences.* 2013;31(13):1491-1502.
77. Hägglund M, Waldén M, Magnusson H, Kristenson K, Bengtsson H, Ekstrand J. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med.* 2013;47(12):738-742.
78. Hirata I, Jr. Editorial: Title IX and the female athlete. *Journal of the American College Health Association.* 1975;24(2):61-62.
79. Parsons JL, Coen SE, Bekker S. Anterior cruciate ligament injury: towards a gendered environmental approach. *British Journal of Sports Medicine.* 2021:bjsports-2020-103173.
80. Ezzat AM, Brussoni M, Mâsse LC, Emery CA. Effect of Anterior Cruciate Ligament Rupture on Physical Activity, Sports Participation, Patient-Reported Health Outcomes, and Physical Function in Young Female Athletes. *The American journal of sports medicine.* 2021;49(6):1460-1469.